

# BIOCOMPATIBILITY OF ACELLULAR DERMAL MATRIX GRAFT EVALUATED IN CULTURE OF MURINE MACROPHAGES

## *AValiação da Biocompatibilidade da Matriz Dérmica Acelular em Cultura de Células*

Ana Paula VENDRAMINI<sup>1</sup>, Rafaela Fernanda MELO<sup>2</sup>, Rosemary Adriana Chiéríci MARCANTONIO<sup>3</sup>, Iracilda Zepone CARLOS<sup>4</sup>

1- DDS, Graduate student, UNESP - Araraquara Pharmaceutical Science School.

2- DDS, Graduate student, UNESP - Araraquara Dental School.

3- DDS, PhD, Associate Professor, UNESP - Araraquara Dental School.

4- DDS, PhD, Associate Professor, UNESP - Araraquara Pharmaceutical Science School.

**Corresponding address:** Rosemary Adriana Chiéríci Marcantonio - Faculdade de Odontologia de Araraquara – UNESP  
Rua Humaitá, 1680, CEP: 14801-903 - Telefone: (16)330166431, fax: (16)33016433 - Email: [adriana@foar.unesp.br](mailto:adriana@foar.unesp.br)

**Received:** November 18, 2004 - **Modification:** October 20, 2005 - **Accepted:** February 14, 2006

### ABSTRACT

The acellular dermal matrix allograft has been used as an alternative to autogenous palatal mucosal graft. The aim of this study was the evaluation of the biocompatibility of an acellular dermal matrix (AlloDerm®) in culture of macrophages. For hydrogen peroxidase determination we used the method of Pick & Kesari, and the Griess method for nitric oxide determination. Statistical analysis showed no significant difference ( $p \leq 0,05$ ) in the release of nitric oxide and hydrogen peroxide by the macrophages exposed to acellular dermal matrix and the negative control. The results suggest that acellular dermal matrix did not activate the cell inflammatory response.

**Uniterms:** Dermal allograft; Macrophages; Nitric oxide; Hydrogen peroxide.

### RESUMO

A matrix dérmica acelular tem sido utilizada como alternativa para a substituição de enxerto gengival autógeno. O objetivo deste estudo foi avaliar a biocompatibilidade em cultura de células de macrófagos da matriz dérmica acelular (AlloDerm®). Foram utilizados os métodos de Pick & Kesari, para a determinação da presença de peróxido de hidrogênio ( $H_2O_2$ ) e de Griess para a determinação de ácido nítrico (NO). Não houve diferença estatisticamente significativa ( $p \leq 0,05$ ) no aumento da presença de NO e  $H_2O_2$  quando macrófagos foram expostos na presença da matrix dérmica acelular quando comparado com o controle negativo. Pode-se concluir que a matrix dérmica acelular é biocompatível aos tecidos humanos.

**Unitermos:** Matriz dérmica; Macrófagos; Ácido nítrico; Peróxido de hidrogênio.

### INTRODUCTION

Several techniques have been developed to solve gingival aesthetic problems. The techniques can be grouped as pediculated grafts and autogenous grafts (epithelial free grafts and connective tissue graft)<sup>16,22</sup>. Aiming at reducing the problems with these treatments, new techniques and materials have been developed<sup>6,9,18-22,24,26,31</sup>.

A recent material developed for soft tissue grafting was the acellular dermal matrix (AlloDerm®, Lifecore Biomedical, Oral Restorative Division, Chaska,MN)<sup>1,6,25,30,32</sup>. This allograft

material has advantages to reduce surgical complications, decrease donor site discomfort and improve esthetic tissue. This graft is obtained from an allograft donor skin and produced by a carefully controlled process that removes the epidermis and dermis cells without altering the extracellular matrix structure, reducing the possible immune response and the transmission of diseases.

The need for biocompatible materials implies the necessity of in vitro toxicity tests, animal experimentation usage tests and clinical studies in humans<sup>23</sup>. The biological system used in in vitro cytotoxicity testing of some materials

are cells in culture. Thus, culture of macrophage cells can be used to evaluate the biocompatibility of materials. These cells participate in many aspects of host defense, inflammation and immunity, partly through their ability to undergo adaptive responses to the conditions or stimuli that prevail at sites to which they have been attracted<sup>2</sup>.

Among the numerous secretory products of macrophages there are two groups of inorganic compounds with a high degree of chemical reactivity: the ROI (including super oxide, hydrogen peroxide, and in some populations of mononuclear phagocytes, the products of myeloperoxidase), and the RNI including nitrite ( $\text{NO}_2^-$ ) and highly related reactive oxides such as nitric oxide and nitrogen dioxide<sup>3</sup>. Macrophage-derived RNI are of interest for at least three reasons: their production is under strict immunological control<sup>27,28</sup>, they are synthesized by enzymes novel to mammalian biochemistry, which have not been yet well characterized<sup>13</sup>, and they appear to play an important role in some of the carcinogenic<sup>13,15</sup>, antitumor<sup>12,29</sup> and antimicrobial<sup>7</sup> actions of the activated macrophage.

Although Alloderm® is tested before being released for sale; we assume that it is important to evaluate the biocompatibility in macrophages culture.

## MATERIAL AND METHODS

This study was approved by the Institutional Committee on Animal Research, School of Pharmaceutical Science.

### Animals

The Animals Laboratory of the School of Pharmaceutical Science, UNESP (State University of São Paulo) Araraquara, SP, BRAZIL supplied six-week old male Swiss mice weighing 18 to 25 g.

### Acellular dermal matrix samples

Samples of acellular dermal matrix (ADM), commercially known as AlloDerm®, measuring 4x4 mm, from different batches were obtained at the Department of Periodontology of Araraquara Dental School UNESP-Araraquara, São Paulo, Brazil.

### Cell macrophage

Mice were injected i.p. 3 to 4 days before harvesting with 3 ml of thioglycollate broth. Macrophages were obtained after killing the mice with chloroform, and the peritoneum was exposed using sterile scissors. Saline solution (0.85% NaCl) was introduced into the peritoneum and after digital massage, the suspension was removed by aspiration. This suspension of peritoneal cells and saline was placed in a Neubauer chamber and counted in order to obtain the ideal concentration for each test. The adherent cells of PEC were re-suspended in RPMI-1640 medium at concentration of  $5 \times 10^6$  cells/ml, and 100  $\mu$ l of this suspension was added to each well tissue dish along with 100  $\mu$ l of a 10.0  $\mu$ g/ml (1.0  $\mu$ g) E. coli O111B lipopolysaccharide (LPS) solution was used as positive control, or RPMI-1640 medium

as negative control. The cells were incubated for 24 h before stimulation, and 50  $\mu$ l aliquots of culture supernatant were mixed with 50  $\mu$ l of Griess reagent (1% w/v sulphonylamide, 0.1% w/v naphthylethylenediamine and 3%  $\text{H}_3\text{PO}_4$ ), incubated at room temperature for 10 min, and the color reaction was determined at 450 nm with a Multiskan Ascent ELISA reader (Labsystems, Helsinki, Finland) Supernatants from quadruplicate cultures were assayed in four experiments and reported as the mean  $\text{NO}_2$  concentration  $\pm$ SD.

### $\text{NO}_2$ release

Nitrite concentration in the medium was measured by a micro plate assay method<sup>5</sup>. The Alloderm® was incubated with cell suspension in a concentration of  $5 \times 10^6$  cells / ml, at 37°C and atmosphere with  $\text{CO}_2$  5%. After 24 hs, 50  $\mu$ l aliquots were removed from conditioned medium and incubated with an equal volume of Griess reagent (1% sulfanilamide, 0.1% naphthylethylene diamine dihydrochloride, 0.25%  $\text{H}_3\text{PO}_4$ ) at room temperature for 10 minutes. The absorbance at 550 nm was determined in a micro plate reader. The tests were made in quadruplicate and the results were expressed in micromoles of NO /  $5 \times 10^5$  peritoneal cells, from a standard curve established in each test, constituted of known molar concentrations of NO in RPMI – 1640 medium.

### $\text{H}_2\text{O}_2$ release

The method depends on the determination of hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) release in the culture of peritoneal macrophages from Swiss mice. Suspensions of peritoneal cells were performed using a concentration of  $2 \times 10^6$  cells/ml in a solution of phenol red, containing 140 mM NaCl, 10 mM potassium phosphate, pH 7.0; 5.5 mM dextrose; 0.56 mM phenol red and type II horseradish peroxidase 0.01 mg/mL (Sigma). Aliquots of 0.1 ml were transferred to culture plates, flat bottomed containing 96 wells (Corning). The acellular dermal matrix or 50  $\mu$ l of Zymosan solution (5 mg/ml, Sigma) were added to each well. The samples were incubated for one hour at 37°C in a 5%  $\text{CO}_2$  atmosphere. After the period of incubation, the reaction was interrupted by addition of 10  $\mu$ l of NaOH 4N. Experiments were done in quadruplicate and the absorbance was determined in an ELISA automatic photometer, with a 620 nm filter. The results were expressed in nanomols of  $\text{H}_2\text{O}_2$  /  $2 \times 10^5$  peritoneal cells, from a standard curve established in each test, constituted of known molar concentrations of  $\text{H}_2\text{O}_2$  in buffered phenol red.

### Statistical Analysis

Comparisons between groups were performed using Student's T test. P values <0.05 were considered statistically significant.


## RESULTS

Figures 1 and 2 illustrate the release of nitric oxide and hydrogen peroxide when macrophages were exposed to


acellular dermal matrix, respectively. Statistical analysis showed no significant difference ( $p \leq 0.05$ ) when compared with negative control. However, the level of release of these mediators in positive control was statistically different ( $p \geq 0.05$ ).

## DISCUSSION

This study evaluated the cell inflammatory response of *in vitro* macrophages in contact with the acellular dermal matrix measuring the release of nitric oxide and hydrogen peroxide. The positive control using zymosan-SIGMA to  $H_2O_2$  release and LPS-SIGMA to NO release showed a high potential of macrophages activation and liberation of  $H_2O_2$  and NO, respectively. The acellular dermal matrix was not able to stimulate macrophage liberation of hydrogen peroxide ( $H_2O_2$ ) and nitric oxide (NO).



**FIGURE 1-** Production of nitric oxide by peritoneal macrophages cultured with medium (C), acellular dermal matrix and lipopolysaccharide ( $1 \times 10^3$  mg/well), by 24 hours of incubation at 37 degree. The results were expressed as the mean  $\pm$  SD of quadruplicate from five determinations



**FIGURE 2-** Production of hydrogen peroxide by peritoneal macrophages cultured with medium, acellular dermal matrix and Zymosan ( $250 \mu\text{g/well}$ ). In the test phenol red was used as indicator of presence of hydrogen peroxide. The results were expressed as the mean  $\pm$  SD of quadruplicate from five determinations

In respect to these results we pointed out that Liversey, et al.<sup>14</sup> (1994) evaluating immunohistochemically the material AlloDerm® did not find any antigen that could develop an immune reaction.

Cellular events were evaluated in some human studies with ADM grafts for root coverage<sup>4,10,11,33</sup>. In 1998, Harris<sup>10</sup>, using acellular dermal matrix graft in treatment of gingival recessions in patients, obtained complete root coverage on two of three defects. In histological analysis, the AlloDerm® had incorporated and became part of the gingival tissue in the area. The same author<sup>11</sup> in 2001 reported a comparative clinical study of root coverage obtained with ADM versus connective tissue. Biopsy of the grafted area revealed elastin fibers. The author stated that the presence of these fibers implied that the ADM was being incorporated into the host tissue.

In a recent study Cummings, et al.<sup>4</sup> (2005) histologically evaluated the acellular dermal matrix graft. The findings of the study showed new fibroblast, vascular elements, and collagen were present throughout the ADM, with retention of the transplanted elastin fibers.

All of these results can be directly related with the laboratory processing of human skin obtained in tissue banks with the removal of all cells without altering the connective tissue structure, composed of type I collagen fibers<sup>14</sup>. In this respect, the AlloDerm® matrix is rigorously controlled by the FDA according to the guidelines of the American Association of Tissue Banks. Apart from the selection of possible donors, serologic and microbiologic exams are also performed to screen for diseases such as AIDS, syphilis, hepatitis, etc. The patented process of preparation and lyophilization removes all cells, preserving the collagen structure of the connective tissue. In the manufactured controls, histological and immunohistochemistry tests are carried out to check the complete removal of all cellular components.

Within the limits of this study the present results show that there was no release of hydrogen peroxide ( $H_2O_2$ ) and nitric oxide (NO), suggesting that the acellular dermal matrix did not activate the cell inflammatory response, although new studies should be accomplished with AlloDerm® use.

## REFERENCES

- 1- Callan DP. Use of acellular dermal matrix allograft material in dental implant treatment. Dent Surg Products. 1996;14-7.
- 2- Chan ED, Winston BW, Uh ST, Wynes MW, Rose DM, Riches DW. Evaluation of the role of mitogen-activated protein kinases in the expressions of inducible nitric oxide synthase by IFN- $\gamma$  and TNF- $\alpha$  in mouse macrophages. J Immunol. 1999;162(1):415-22
- 3- Cotton FA, Wilkinson G. Advanced inorganic chemistry, a comprehensive text, 4<sup>th</sup> ed. New York: Wiley-Interscience; 1980. p. 422-30.
- 4- Cummings LC, Kaldahl WB, Allen EP. Histologic evaluation of autogenous connective tissue and acellular dermal matrix graft in humans. J Periodontol. 2005;76(2):178-86.

- 5- Ding AH, Nathan CF, Stuehr DJ. Release of reactive nitrogen intermediates and reactive oxygen intermediates from mouse peritoneal macrophages. Comparison of activating cytokines and evidence for independent production. *J Immunol.* 1988;141(7):2407-12
- 6- Dodge JD, Henderson R, Greenwell H. Root coverage without a palatal donor site, using an cellular dermal graft. *Periodontal Insights.* 1998;5-8.
- 7- Granger DL, Hibbs JB Jr, Perfect JR, Durack DT. Specific amino acid (L-arginine) requirement for the microbiostatic activity of murine macrophages. *J Clin Invest.* 1988;81(4):1129-36.
- 8- Harris RJ. The connective tissue with partial thickness double pedicle graft: the results of 100 consecutively treated defects. *J Periodontol.* 1994;65(5):448-61.
- 9- Harris RJ. A comparative study of root coverage obtained with guided tissue regeneration utilizing a bioabsorbable membrane versus the connective tissue with partial-thickness double pedicle graft. *J Periodontol.* 1997;68(8): 779-90.
- 10- Harris RJ. Root coverage with a connective tissue with partial thickness double pedicle graft and an acellular dermal matrix graft: A clinical and histological evaluation of a case report. *J Periodontol.* 1998;69(11):1305-11.
- 11- Harris RJ. Gingival augmentation with an acellular dermal matrix: Human histological evaluation of a case – placement of the graft on bone. *Int J Periodontics Restorative Dent.* 2001;21(1):69-75.
- 12- Hibbs JB Jr, Taintor RR, Vavrin Z. Macrophage cytotoxicity: role for L-arginine deiminase and imino nitrogen oxidation to nitrite. *Science.* 1987 Jan 23;235(4787):473-6.
- 13- Iyengar R, Stuehr DJ, Marletta MA. Macrophage synthesis of nitrite, nitrate and N-nitrosamines: precursors and role of burst respiratory. *Proc Natl Acad Sci.* 1987;84(18):6369-73.
- 14- Livesey S, et al. An acellular dermal transplant processed from human allograft skin retains normal extracellular matrix components and ultrastructural characteristics (poster). *AATB Conference;* 1994 august.
- 15- Miwa M, Stuehr DJ, Marletta MA, Wishnok, JS, Tannenbaum SR. Nitrosation of amines by stimulated macrophages. *Carcinogenesis.* 1987;8(7):955-8.
- 16- Miller PD Jr. Root coverage using the free soft tissue autograft following citric acid application III. A successful and predictable procedure in areas of deep-wide recession. *Int J Periodontics Restorative Dent* 1985;5(2):14-37.
- 17- Pini Prato G, Tinti C, Vincenzi G, Masnani C, Cortellini P, Clauser C. Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal gingival recession. *J Periodontol.* 1992;63(11):919-28.
- 18- Pini Prato G, Clauser C, Cortellini P, Tinti C, Vincenzi G, Pagliaro U. Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal recessions. A 4-year follow-up study. *J Periodontol.* 1996;67(11):1216-23
- 19- Pini Prato GP, Clauser C, Cortellini P. Periodontal plastic and mucogingival surgery. *Periodontol* 2000 1995;9:90-105.
- 20- Rachlin G, Koubi G, Dejou J, Franquin JC. The use of a resorbable membrane in mucogingival surgery. Case series. *J Periodontol.* 1996;67(6):621-6.
- 21- Ricci G, Silvestri M, Tinti C, Rasperini G. A clinical/ statistical comparison between the subpedicle connective tissue graft method and the guided tissue regeneration technique in root coverage. *Int J Periodontics Restorative Dent.* 1996;16(6):539-45.
- 22- Sbordone L, Ramaglia L, Spagnuolo G, De Luca M. A comparative study of free gingival auto graft and free gingival and subepithelial; connective tissue grafts. *Periodontal Case Reports.* 1988;10(1):8-12.
- 23- Schmalz G. Use of cell cultures for toxicity testing of dental materials-advantages and limitations. *J Dent.* 1994;22(Suppl 2):S6-11.
- 24- Shieh AT, Wang HL, O'Neal R, Glickman GN, MacNeil RL. Development and clinical evaluation of a root coverage procedure using a collagen barrier membrane. *J Periodontol.* 1997;68(8):770-8.
- 25- Silverstein LH, Callan DP. An acellular dermal matrix allograft substitute for palatal donor tissue. *Postgrad Dent.* 1996;3:14-21.
- 26- Silverstein LH. Fundamentally changing soft tissue grafting. *Dentistry Today.* 1997;16: 1-3.
- 27- Stuehr DJ, Marletta MA. Mammalian nitrate biosynthesis: mouse macrophages produce nitrite and nitrate in response to Escherichia coli lipopolysaccharide. *Proc Natl Acad Sci USA.* 1985 Nov;82(22):7738-42.
- 28- Stuehr DJ, Marletta MA. Induction of nitrite/nitrate synthesis in murine macrophages by BCG infection, lymphokines, or interferon. *J Immunol.* 1987;139(2):518-25.
- 29- Stuehr DJ, Marletta MA. Cytostasis from nitrite, a product of activated macrophages. *Cancer Res.* 1987;47(21):5590-4.
- 30- Tal H. Subgingival acellular dermal matrix allograft for the treatment of gingival recession: A case report. *J Periodontol.* 1999;70:1018-24.
- 31- Vernino AR, Young SK, Tow HD. Histologic evaluation following intraoral use of freeze-dried skin in humans. *Int J Periodontics Restorative Res.* 1986;6(4):57-65.
- 32- Wainwright DJ. Use of an acellular allograft dermal matrix (AlloDerm®) in management of full-thickness burns. *Burns.* 1995;21(4):243-8.
- 33- Wei P, Laurell L, Lingen MW, Geivellis M. Acellular Dermal Matrix allografts to achieve increased attached gingiva. Part 2. A histological comparative study. *J Periodontol.* 2002;73(3):257-65.